#### Class X Mathematics –Standard (041) Sample Question Paper 2019-20

Max. Marks: 80

**Duration : 3 hrs** 


#### General Instructions:

- (i) All the questions are compulsory.
- (ii) The question paper consists of 40 questions divided into 4 sections A, B, C, and D.
- (iii) Section A comprises of 20 questions of 1 mark each. Section B comprises of 6 questions of 2 marks each. Section C comprises of 8 questions of 3 marks each. Section D comprises of 6 questions of 4 marks each.
- (iv) There is no overall choice. However, an internal choice has been provided in two questions of 1 mark each, two questions of 2 marks each, three questions of 3 marks each, and three questions of 4 marks each. You have to attempt only one of the alternatives in all such questions.
- (v) Use of calculators is not permitted.


|      | SECTION A  |   |
|------|--|---|
| Q 1  | • Q 10 are multiple choice questions. Select the most appropriate answer from the  |   |
| give | en options.  |   |
| 1    | The decimal representation of $\frac{11}{2^3 \times 5}$ will<br>a) terminate after 1 decimal place<br>b) terminate after 2 decimal places<br>c) terminate after 3 decimal places<br>d) not terminate | 1 |
|      |  |   |

| 2 | Consider t  | ne following   | frequency of     | listribution o   | f the heights | s of 60 stude | ents of a class | 1 |
|---|---|----------------|------------------|--|---------------|---------------|-----------------|---|
|   | Height (in<br>cm)                                 | 150-155        | 155-160          | 160-165  | 165-170       | 170-175       | 175-180         |   |
|   | No of<br>students                                 | 15             | 13               | 10   | 8             | 9             | 5               |   |
|   | The upper<br>a) 165<br>b) 155<br>c) 160<br>d) 170 |                | nedian clas      | s in the give  | n data is     |               |                 |   |
| 3 | The LCM c   | of smallest tv | wo digit com     | posite numl  | per and sma   | llest compos  | site number is  | 1 |
|   | a) 12<br>b) 4<br>c) 20<br>d) 44                   |                |                  |  |               |               |                 |   |
| 4 | For which   | value(s) of p  | o, will the line | es represen  | ted by the fo | llowing pair  | of linear       | 1 |
|   | equations   | be parallel    |                  | 3x - y - 6x - 2y - 6x - 2x |               |               |                 |   |
|   | a) all r<br>b) 10<br>c) 5/2<br>d) 1/2             | eal values e   | xcept 10         | ο <i>λ</i> – 2 <i>y</i> –  | p = 0         |               |                 |   |
|   |   |                |                  |  |               |               |                 |   |

| 5 | If triangle ABC is right angled at C, then the value of sec (A+B) is<br>a) 0<br>b) 1<br>c) $\frac{2}{\sqrt{3}}$<br>d) not defined  | 1 |
|---|--|---|
| 6 | If $sin\theta + cos\theta = \sqrt{2}cos\theta$ , $(\theta \neq 90^{\circ})$ then the value of $tan\theta$ is<br>a) $\sqrt{2} - 1$<br>b) $\sqrt{2} + 1$<br>c) $\sqrt{2}$<br>d) $-\sqrt{2}$        | 1 |
| 7 | Given that $sin\alpha = \frac{\sqrt{3}}{2}$ and $cos\beta = 0$ , then the value of $\beta - \alpha$ is<br>a) 0°<br>b) 90°<br>c) 60°<br>d) 30°  | 1 |
| 8 | The point which divides the line segment joining the points (8, – 9) and (2, 3) in<br>ratio 1 : 2 internally lies in the<br>a) I quadrant<br>b) II quadrant<br>c) III quadrant<br>d) IV quadrant | 1 |
| 9 | The distance of the point P (-3, -4) from the <i>x</i> -axis (in units) is<br>a) 3<br>b) -3<br>c) 4<br>d) 5  | 1 |




| 14   | Fill the two blanks in the sequence 2,, 26, so that the sequence forms an A.P   | 1 |
|------|---|---|
| 15   | A number is chosen at random from the numbers -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5. Then the probability that square of this number is less than or equal to 1 is               | 1 |
| (Q ′ | I6- Q 20) Answer the following  |   |
| 16   | Write one rational and one irrational number lying between 0.25 and 0.32  | 1 |
| 17   | In the figure, if $\angle ACB = \angle CDA$ , $AC = 6 \text{ cm}$ and $AD = 3 \text{ cm}$ , then find the length of AB  | 1 |
| 18   | D<br>If the angle between two tangents drawn from an external point 'P' to a circle of radius 'r'<br>and centre O is 60 <sup>0</sup> , then find the length of OP.<br><b>OR</b> | 1 |
|      | If the radii of two concentric circles are 4 cm and 5 cm, then find the length of each chord of one circle which is tangent to the other circle.                                |   |
| 19   | If the first three terms of an A.P are b, c and 2b, then find the ratio of b and c  | 1 |
| 20   | Find the value(s) of k for which the quadratic equation $x^2 + 2\sqrt{2}kx + 18 = 0$ has equal roots  | 1 |
|      | <u>Section – B</u>  |   |
| 21   | Find the number of natural numbers between 102 and 998 which are divisible by 2 and 5 both.   | 2 |
| 22   | Prove that the rectangle circumscribing a circle is a square.   | 2 |



| 25 | Jayanti throws a pair of dice and records the product of the numbers appearing on the  | 2 |
|----|--|---|
|    | dice. Pihu throws 1 dice and records the squares the number that appears on it. Who  |   |
|    | has the better chance of getting the number 36? Justify?   |   |
|    | OR   |   |
|    | <ul> <li>An integer is chosen between 70 and 100, Find the probability that it is</li> <li>(a) a prime number</li> <li>(b) divisible by 7</li> </ul>   |   |
|    |  |   |
| 26 | Isha is 10 years old girl. On the result day, Isha and her father Suresh were very happy<br>as she got first position in the class. While coming back to their home, Isha asked for a<br>treat from her father as a reward for her success. They went to a juice shop and asked<br>for two glasses of juice. | 2 |
|    | Aisha, a juice seller, was serving juice to her customers in two types of glasses.<br>Both the glasses had inner radius 3cm. The height of both the glasses was 10cm.  |   |
|    |  |   |
|    | First type: A Glass with hemispherical raised bottom.  |   |
|    |  |   |
|    | Second type: A glass with conical raised bottom of height 1.5 cm.  |   |
|    | Isha insisted to have the juice in first type of glass and her father decided to have the juice in second type of glass. Out of the two, Isha or her father Suresh, who got more quantity of juice to drink and by how much?   |   |
|    |  |   |
|    | Section C  |   |
|    | Section C  |   |
| 27 | Given that $\sqrt{5}$ is irrational, prove that $2\sqrt{5} - 3$ is an irrational number.   | 3 |
|    | OR   |   |
|    | OK   |   |
|    | If HCF of 144 and 180 is expressed in the form 13m-16. Find the value of m.  |   |

| 28 | If the sum of first m terms of an AP is the same as the sum of its first n terms, show that the sum of its first (m+n) terms is zero.   | 3 |
|----|---|---|
| 20 | In the figure ABCDE is a pentagen with BEIICD and BCIIDE, BC is perpendicular to  | 3 |
| 29 | In the figure, ABCDE is a pentagon with BE  CD and BC  DE. BC is perpendicular to CD. AB= 5cm, AE=5cm, BE= 7cm, BC= x-y and CD= x+y. If the perimeter of ABCDE is 27cm. find the value of x and y, given x, $y \neq 0$ .  | 3 |
|    | OR<br>Solve the following system of equations:  |   |
|    | $\frac{21}{x} + \frac{47}{y} = 110$   |   |
|    | $\frac{47}{x} + \frac{21}{y} = 162,  x, y \neq 0$   |   |
| 30 | Obtain all the zeros of the polynomial $x^4+4x^3-2x^2-20x-15$ , if two of its zeroes are $\sqrt{5}$ and $-\sqrt{5}$ .   | 3 |
| 31 | Two friends Seema and Aditya work in the same office at Delhi. In the Christmas vacations, both decided to go to their hometowns represented by Town A and Town B respectively in the figure given below. Town A and Town B are connected by trains from the same station C (in the given figure)in Delhi.Based on the given situation, answer the following questions: | 3 |

|    | Town A<br>6<br>6<br>5<br>Station C  |   |
|----|---|---|
|    | • • • • • • • • • • • • • • • • • • •   |   |
|    | <ul> <li>(i) Who will travel more distance, Seema or Aditya, to reach to their hometown?</li> <li>(ii) Seema and Aditya planned to meet at a location D situated at a point D represented by the mid-point of the line joining the points represented by the point D.</li> <li>(iii) Find the area of the triangle formed by joining the points represented by A, B and C.</li> </ul> |   |
| 32 | If sin $\theta$ + cos $\theta = \sqrt{3}$ , then prove that tan $\theta$ + cot $\theta$ =1  | 3 |
|    | OR<br>Evaluate:<br>$\frac{\cos^{2}(45^{\circ}+\theta) + \cos^{2}(45^{\circ}-\theta)}{\tan(60^{\circ}+\theta) \times \tan(30^{\circ}-\theta)} + (\cot 30^{\circ} + \sin 90^{\circ}) \times (\tan 60^{\circ} - \sec 0^{\circ})$   |   |
| 33 | Sides of a right triangular field are 25m, 24m and 7m. At the three corners of the field, a cow, a buffalo and a horse are tied separately with ropes of 3.5 m each to graze in the field. Find the area of the field that cannot be grazed by these animals.   |   |



| 36       | Prove that sides in dis  |  | •                         | rallel to one<br>other two sig   |   | 0                    |  |              | 4 |
|----------|--|--|---------------------------|--|---|----------------------|--|--------------|---|
| 37       |  |  |                           | ) km at a un<br>nutes less f   | -   |                      | -  |              |   |
|          | Solve the f $\frac{1}{x} = \frac{1}{x}$                              | ollowing equation $\frac{1}{x-2} = 3, x$                                   |                           |  |   |                      |  |              |   |
| 38       | lower and  | upper ends   | s as 20 m a               | a frustum of<br>and 50 m re<br>ate of Rs. 70                               | spectively.   | Find the co          | ost of petro   | l which can  |   |
|          |  |  |                           | _  |   |                      |  |              |   |
|          |  |  |                           | C  | R   |                      |  |              |   |
|          |  | ond which i  |                           | <b>C</b><br>f 15km/hou<br>g and 44m w                                      | r through a   |                      |  |              |   |
| 39       | cuboidal po<br>pond rise b<br>The angle<br>30 second                 | ond which i<br>by 21cm?<br>of elevatio<br>s, the angl                      | n of an air<br>e of eleva | f 15km/hou   | r through a<br>vide. In what<br>a point on th<br>es 30°. If the                         | time will the ground | ne level of v  | water in the | 4 |
|          | cuboidal po<br>pond rise to<br>The angle<br>30 second<br>height of 3 | ond which in<br>by 21cm?<br>of elevations, the angle<br>$000\sqrt{3}$ m, f | n of an air<br>e of eleva | f 15km/hou<br>g and 44m w<br>plane from a<br>tion become<br>eed of the air | r through a<br>ride. In what<br>a point on th<br>es 30 <sup>0</sup> . If the<br>rplane. | time will the ground | ne level of v<br>is 60 <sup>0</sup> . Afte<br>is flying at | water in the | 4 |
| 39<br>40 | cuboidal po<br>pond rise to<br>The angle<br>30 second<br>height of 3 | ond which in<br>by 21cm?<br>of elevations, the angle<br>$000\sqrt{3}$ m, f | n of an air<br>e of eleva | f 15km/hou<br>g and 44m w<br>plane from a<br>tion become                   | r through a<br>ride. In what<br>a point on th<br>es 30 <sup>0</sup> . If the<br>rplane. | time will the ground | ne level of v<br>is 60 <sup>0</sup> . Afte<br>is flying at | water in the | 4 |